2025 Southeast Alaska Herring Spawn-On-Kelp Pound Fishery Management Plan

by Aaron Dupuis Scott Forbes Bo Meredith and Katie Taylor

February 2025

Alaska Department of Fish and Game



Division of Commercial Fisheries

Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figures or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative		all standard mathematical	
deciliter	dL	Code	AAC	signs, symbols and	
gram	g	all commonly accepted		abbreviations	
hectare	ha	abbreviations	e.g., Mr., Mrs.,	alternate hypothesis	H _A
kilogram	kg		AM, PM, etc.	base of natural logarithm	е
kilometer	km	all commonly accepted		catch per unit effort	CPUE
liter	L	professional titles	e.g., Dr., Ph.D.,	coefficient of variation	CV
meter	m		R.N., etc.	common test statistics	(F, t, χ^2 , etc.)
milliliter	mL	at	a	confidence interval	CI
millimeter	mm	compass directions:		correlation coefficient	
		east	E	(multiple)	R
Weights and measures (English)		north	Ν	correlation coefficient	
cubic feet per second	ft ³ /s	south	S	(simple)	r
foot	ft	west	W	covariance	cov
gallon	gal	copyright	©	degree (angular)	0
inch	in	corporate suffixes:		degrees of freedom	df
mile	mi	Company	Co.	expected value	Ε
nautical mile	nmi	Corporation	Corp.	greater than	>
ounce	ΟZ	Incorporated	Inc.	greater than or equal to	≥
pound	lb	Limited	Ltd.	harvest per unit effort	HPUE
quart	qt	District of Columbia	D.C.	less than	<
yard	yd	et alii (and others)	et al.	less than or equal to	\leq
-	2	et cetera (and so forth)	etc.	logarithm (natural)	ln
Time and temperature		exempli gratia		logarithm (base 10)	log
day	d	(for example)	e.g.	logarithm (specify base)	log ₂ , etc.
degrees Celsius	°C	Federal Information		minute (angular)	,
degrees Fahrenheit	°F	Code	FIC	not significant	NS
degrees kelvin	Κ	id est (that is)	i.e.	null hypothesis	Ho
hour	h	latitude or longitude	lat or long	percent	%
minute	min	monetary symbols		probability	Р
second	s	(U.S.)	\$, ¢	probability of a type I error	
		months (tables and		(rejection of the null	
Physics and chemistry		figures): first three		hypothesis when true)	α
all atomic symbols		letters	Jan,,Dec	probability of a type II error	
alternating current	AC	registered trademark	®	(acceptance of the null	
ampere	А	trademark	тм	hypothesis when false)	β
calorie	cal	United States		second (angular)	"
direct current	DC	(adjective)	U.S.	standard deviation	SD
hertz	Hz	United States of		standard error	SE
horsepower	hp	America (noun)	USA	variance	
hydrogen ion activity	pH	U.S.C.	United States	population	Var
(negative log of)	-		Code	sample	var
parts per million	ppm	U.S. state	use two-letter	*	
parts per thousand	ppt,		abbreviations		
- •	%		(e.g., AK, WA)		
volts	V				
watts	W				

REGIONAL INFORMATION REPORT NO. 1J25-06

2025 SOUTHEAST ALASKA HERRING SPAWN-ON-KELP POUND FISHERY MANAGEMENT PLAN

by

Aaron Dupuis Alaska Department of Fish and Game, Division of Commercial Fisheries, Sitka

Scott Forbes Alaska Department of Fish and Game, Division of Commercial Fisheries, Douglas

Bo Meredith Alaska Department of Fish and Game, Division of Commercial Fisheries, Ketchikan

and

Katie Taylor Alaska Department of Fish and Game, Division of Commercial Fisheries, Petersburg

> Alaska Department of Fish and Game Division of Commercial Fisheries, Publications Section 802 3rd Street Douglas, AK 99824-5412

> > February 2025

The Regional Information Report Series was established in 1987 and was redefined in 2007 to meet the Division of Commercial Fisheries regional need for publishing and archiving information such as area management plans, budgetary information, staff comments and opinions to Alaska Board of Fisheries proposals, interim or preliminary data and grant agency reports, special meeting or minor workshop results and other regional information not generally reported elsewhere. Reports in this series may contain raw data and preliminary results. Reports in this series receive varying degrees of regional, biometric and editorial review; information in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author or the Division of Commercial Fisheries if in doubt of the level of review or preliminary nature of the data reported. Regional Information Reports are available through the Alaska State Library and on the Internet at: http://www.adfg.alaska.gov/sf/publications/.

Product names used in this publication are included for completeness and do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

Aaron Dupuis Alaska Department of Fish and Game, Division of Commercial Fisheries, 304 Lake Street. Room 103, Sitka, AK 99835, USA

Scott Forbes Alaska Department of Fish and Game, Division of Commercial Fisheries, 802 3rd Street, Douglas, AK 99824, USA

Bo Meredith Alaska Department of Fish and Game, Division of Commercial Fisheries, 2030 Sea level Drive Suite 205, Ketchikan, AK 99901, USA

and

Katie Taylor Alaska Department of Fish and Game, Division of Commercial Fisheries, 16 Sing Lee Alley, Petersburg, AK 99833, USA

This document should be cited as follows:

Dupuis, A., S. Forbes, B. Meredith, and K. Taylor. 2025. 2025 Southeast Alaska herring spawn-on-kelp pound fishery management plan. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 1J25-06, Douglas.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write: ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526 U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203 Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers: (VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648, (Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact: ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2517

TABLE OF CONTENTS

Page

LIST OF TABLES	ii
LIST OF FIGURES	ii
ABSTRACT	1
INTRODUCTION	1
HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE	2
Methods of Forecasting Herring Biomass Craig/Klawock (Section 3-B) Ernest Sound (District 7) Tenakee Inlet (Section 12-A) Hoonah Sound (Section 13-C)	
CALENDAR OF EVENTS	6
REGULATIONS	6
General Spawn-On-Kelp Regulations Placement and Release of Herring in Pounds	
Post-Harvest Requirements	6
Harvest and Production	6
Requirements for Buyers Closure of some pound types Other Regulations	7
HARVEST AND ALLOCATION OF KELP FOR 2025	7
Section 3-B (Craig/Klawock)	7
FISHERY CONDUCT AND MANAGEMENT	8
Open Waters Closed Waters	
OTHER AGENCY REQUIREMENTS	9
Alaska Department of Natural Resources U.S. Forest Service U.S. National Marine Fisheries Service U.S. Coast Guard	
LIST OF CONTACTS	11
TABLES AND FIGURES	

LIST OF TABLES

Table		Page
1.	Craig/Klawock herring stock size and fishery summary, 1990–2024.	14
2.	Craig/Klawock herring spawn-on-kelp (SOK) fishery summary, 2010–2024.	15
3.	Craig and Ernest Sound herring spa Fisheries Scientist 1wning dates comparison, 2010–2024	16
4.	Ernest Sound herring stock and fishery summary, 1969–2024.	17
5.	Ernest Sound spawn-on-kelp (SOK) fishery summary, 2004, 2008, 2009, 2011, 2013, and 2014	19
6.	Tenakee Inlet herring stock and fishery summary, 1978–2024.	
7.	Tenakee Inlet herring Spawn-on-kelp (SOK) fishery summary, 2003–2005, 2009, and 2014	
8.	Hoonah Sound and Tenakee Inlet herring spawn dates comparison, 2003–2024.	
9.	Hoonah Sound herring spawning stock and spawn-on-kelp (SOK) fishery summary, 1971–2015	
10.	Hoonah Sound herring spawn-on-kelp (SOK) detailed fishery summary, 2003–2013.	
		-

LIST OF FIGURES

Figure		Page
1.	Craig/Klawock (Section 3-B) herring SOK fishery open area.	26
2.	Diagram of a herring pound showing 2 alternative methods of marking herring pounds.	27
3.	U.S. Coast Guard requirements for marking pounds	

ABSTRACT

This management plan provides an overview of the management approach, permit requirements, and regulations for the 2025 herring spawn-on-kelp (SOK) pound fisheries in Southeast Alaska. A herring SOK pound fishery will only occur in the Section 3-B (Craig/Klawock) area. The forecasted mature spawning biomass in Section 3-B is 30,182 tons of herring for the 2024–2025 season. This biomass estimate allows for a 20% harvest rate and a combined guideline harvest level (GHL) of 6,036 tons for the winter food and bait and the SOK fishery. The minimum GHL available to the Section 3-B SOK fishery is 2,414 tons, allowing for full kelp allocation.

Keywords: Pacific herring, Clupea pallasii, herring pound, Macrocystis kelp, allocation, management plan, spawn-on-kelp

INTRODUCTION

This plan provides an overview of the 2025 management approach, permit requirements, and regulations for Pacific herring *Clupea pallasii* harvest in the Southeast Alaska spawn-on-kelp (SOK) fisheries. 5 AAC 27.185 *Management plan for herring spawn on kelp in pounds fisheries in Sections 3-B, 12-A, and 13-C, and District 7* establishes the regulatory framework for the Southeast Alaska SOK fisheries and provides for fisheries in Sections 3-B (Craig/Klawock), 12-A (Tenakee Inlet), 13-C (Hoonah Sound), and in District 7 (Ernest Sound).

A *closed-pound fishery* involves capturing sexually mature herring and releasing them into a net impoundment in which kelp is suspended. The herring are released from the pound after they spawn on the suspended kelp and the kelp with eggs attached is then sold. An *open-pound fishery* involves suspending kelp from a floating frame structure in an area where herring are spawning. The herring are not impounded but instead are allowed to naturally spawn on the suspended kelp. The kelp blades with eggs are removed from the water then sold. In Southeast Alaska herring SOK fisheries, a closed or an open pound may be operated by one or more Commercial Fisheries Entry Commission (CFEC) permit holders.

The 2024–2025 season (October 1–September 30) herring guideline harvest level (GHL) for the Craig/Klawock area stock is 6,036 tons of herring. Forty percent (2,414 tons) is allocated to the SOK fishery plus any unharvested portion of the winter food and bait quota. The current GHL for the Craig/Klawock SOK fishery corresponds to an allocation range of *1,500 or more* tons of kelp; this allocation range provides for the maximum allowable number of kelp blades for each permit holder by pound type.

No fishery will occur in Ernest Sound during the 2024–2025 season. A forecast was not generated due to the small amount of herring spawn observed in 2024. In 2025, Ernest Sound will be monitored throughout the duration of historical spawn timing. If enough herring spawn is observed and funding allows, a spawn deposition survey may be conducted.

No fishery will occur in Hoonah Sound during the 2024–2025 season. In 2024, no herring or herring spawn were observed, therefore no forecast was generated for the 2024–2025 season. Future assessments and fisheries are dependent on available funding and documentation of herring spawn.

No fishery will occur in Tenakee Inlet during the 2024–2025 season. In 2024, approximately 10.6 nmi of herring spawn was documented for this stock (4.8 nmi from inside Tenakee Inlet south to Basket Bay and 5.8 nmi from Basket Bay south to Lull Point). A spawn deposition survey was conducted in 2024, but no forecast was generated for the 2024–2025 season. In 2025, the stock will be monitored throughout historical spawn timing. If enough herring spawn is observed and funding allows, a spawn deposition survey may be conducted.

HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE

METHODS OF FORECASTING HERRING BIOMASS

The age-structured assessment (ASA) models (also called statistical catch-at-age models) are used to forecast herring biomass for several of the larger stocks in Southeast Alaska, including Craig/Klawock. These models primarily rely on a time series of estimated total egg deposition, spawner-age composition, catch-age composition, weight at age, and harvest. Estimates of fecundity are also included in the models. The models estimate recruitment, maturation, and natural mortality and apply them to an estimate of spawning biomass from a given year to forecast mature biomass for the following year. The ASA model was an important development because gains in herring biomass due to recruitment, growth, and maturity are often not equal to the loss of biomass due to natural mortality, as was assumed when using the prior year spawn deposition estimate for forecasting biomass.

The biomass accounting (BA) method of forecasting is a simpler alternative to the ASA method, and it is used for areas with historical datasets that are shorter or less complete, or years with insufficient time to process data or an assessment. The BA method is used to forecast the prefishery mature biomass; it has been used to establish the fishery GHL in Hoonah Sound (most recent forecast in 2016), Ernest Sound (the most recent forecast in 2017), and was used to forecast the 2023 mature biomass of Craig herring. The BA method uses only the current year's estimate of egg deposition, the current age compositions of the spawning and commercially caught biomass, weight at age, and fecundity estimates to project the following year's return of mature herring biomass. The Hoonah Sound forecast uses the estimated survival, estimated maturity, and fecundity relationship derived from the ASA model for the nearby Seymour herring stock. A median historical level of recruitment of age-3 herring specific to Hoonah Sound is also applied to forecast biomass. The Ernest Sound forecast uses the estimated survival, estimated maturity, and fecundity relationship derived from the ASA model for the nearby Craig herring stock. A median historical recruitment of age-3 herring specific to Ernest Sound is also applied to forecast biomass. The BA model was used to forecast the 2023 mature biomass of the Craig herring stock for 3 reasons: biometric time constraints, near certainty that the forecast would be above threshold, and a high likelihood that the forecast would result in GHLs that would not constrain the bait or pound fisheries (taking into account historical bait harvest levels and maximum blade allocation). The ASA model was used to forecast the 2024 and 2025 mature biomass of herring in Craig.

Once a forecast of the season's biomass is calculated, a variable harvest rate formula is applied; that formula determines a harvest rate of 10-20% of the forecast of mature spawning biomass. When the spawning biomass forecast for an area equals the threshold, the exploitation rate is 10% of the estimated spawning biomass. For each incremental increase in the spawning biomass equal to the threshold, the exploitation rate increases by 2%.

CRAIG/KLAWOCK (SECTION 3-B)

Herring fisheries have occurred in Section 3-B since the 1960s. The first commercial harvest was in the form of a wild SOK fishery in the Craig/Klawock area beginning in the early 1960s that ended in 1967. A small winter bait herring fishery began in 1965. Initial harvests were small, less than 5 tons through 1973, and a formal GHL was never established. The use of hydroacoustic surveys began in 1973 and during the subsequent years the winter bait harvest increased with an average harvest of 1,147 tons from 1973 through 1991. The Section 3-B SOK fishery was

established by the board in 1992. The GHL for the Section 3-B stock was allocated between the winter food and bait fishery and the herring SOK fishery. When the fishery was created in 1992, the GHL allocation was 85% for the winter food and bait fishery and 15% for the SOK fishery. In 1998, the allocation was modified to 60% to the winter food and bait fishery with the remaining 40% and any unharvested portion of the food and bait fishery allocated to the herring SOK fishery. Since 1992, the Craig/Klawock herring GHL has averaged 3,503 tons, ranging from a low of 626 tons in 2000 to a high of 19,456 tons in 2021 (Table 1). The herring SOK fishery effort, harvest, spawning dates, fishery dates, and product values are summarized in Table 2.

Annual harvest levels are based on a formula that allows for higher harvest rates as the herring population increases relative to the threshold level. No harvest is allowed if the biomass estimate for the stock is less than the threshold level. The established threshold level for the Craig/Klawock stock is 5,000 tons.

The forecasted mature spawning biomass for the Craig/Klawock stock for 2024–2025 is 30,182 tons of herring. The 2024–2025 Craig/Klawock forecast was derived using the ASA model. This forecast allows for a 20% harvest rate, equaling a total GHL of 6,036 tons for the winter food and bait and SOK fisheries. Therefore, the GHL is 3,622 tons to the winter food and bait fishery and 2,414 tons to the SOK fishery. The Alaska Department of Fish and Game (ADF&G or department) anticipates the dominant age classes in the forecast (by numbers) to be age-5 (43%) and age-8+ (20%) totaling 63% of the mature population.

The 2024–2025 Craig/Klawock winter food and bait fishery will close by regulation on February 28, 2025. All unharvested winter food and bait quota will be added to the GHL for the SOK fishery. The initial allocation of 2,414 tons for the Section 3-B SOK GHL falls within the allocation range of *1,500 tons or more*, which corresponds to the fishery's maximum allowable number of kelp blades per permit holder by pound type.

Herring spawning normally occurs in the Craig/Klawock area between mid-March and early April. The earliest spawning observed since the mid-1970s was March 9 and the latest date of initial spawning was April 9 (Table 3). In 2024, the main spawn event in the Craig/Klawock area began to escalate on March 21, 2024.

ERNEST SOUND (DISTRICT 7)

The Ernest Sound SOK pound fishery was created in January 2003 by the Board of Fisheries (BOF). Additionally, the BOF created a herring bait pound fishery that is allocated 10% of the area's GHL. The bait pound fishery is similar to other herring fisheries in that its allocation is based upon the GHL remaining after the herring food and bait fisheries occur. Any remaining GHL from the winter food and bait fishery or the bait pound fishery after March 15 is allocated to the SOK fishery. SOK fisheries in Ernest Sound have occurred intermittently and at various levels of effort and harvest since the first fishery occurred in 2004 (Table 4). From 2004 through 2024, there have been 6 years in which SOK GHLs were allotted. Effort varied from zero participants in 2011 to 129 in 2014 (Table 5).

The Ernest Sound herring stock has a threshold level of 2,500 tons. A forecast was not developed for 2025 because no spawn deposition survey was conducted. Therefore, no commercial herring fisheries will occur in Ernest Sound during the 2024–2025 season. Historical spawning biomass, forecast, GHLs, spawning dates, harvest, and fishery dates are summarized in Tables 3, 4, and 5.

ADF&G plans to monitor herring spawn in 2025. Aerial and skiff surveys for the past few years have indicated an increasing trend in egg deposition. If herring spawn is substantial, samples of spawning herring may be collected and a spawn deposition survey may be conducted.

TENAKEE INLET (SECTION 12-A)

The Tenakee Inlet herring stock has been the target of a winter food and bait fishery since the 1978–1979 season. During seasons that the estimated spawning biomass was above the 3,000-ton threshold, the GHL ranged from a low of 200 tons in the initial season to a peak of 1,700 tons in the 1985–1986 season (Table 5). Regulations adopted by the BOF in January 2003 provide for a SOK fishery in Tenakee Inlet if sufficient GHL remains at the close of the winter food and bait fishery. The SOK fishery occurred for the first time in April 2003. Summaries of the Tenakee Inlet SOK fisheries are presented in Table 6.

ADF&G has conducted aerial surveys in Tenakee Inlet since the early 1970s, documenting the total miles of spawn each season to provide an indication of herring stock size or biomass. Aerial surveys were supplemented with hydroacoustic surveys from 1979 through 1986. Spawn deposition surveys began in 1987 as the most reliable and accurate means to assess the spawning biomass, but have not been conducted in Tenakee Inlet since 2015.

The Tenakee Inlet stock includes herring that spawn within the inlet itself, along the Chichagof and Catherine Islands shorelines in Chatham Strait between South Passage Point and Point Lull, and in the entrance of Peril Strait east of the longitude of Point Moses. This spawning stock has exhibited cycles of abundance. After a decade of fisheries, the stock declined below threshold in the early 1990s, and no fisheries took place until the 1996–1997 season. Good recruitment led to nearly a decade of harvestable surplus until the forecasted biomass again declined below threshold in 2006. Aerial spawn surveys and spawn deposition dive surveys conducted in 2008 indicated a significant increase in spawning biomass to the levels seen in 1997–1999. However, surveys conducted since 2009 again indicate a decreasing trend in mature spawning biomass. The herring spawn mileage observed in 2014, 2015, and 2017 was approximately 2.0 nmi each year, but no herring spawn was documented within the inlet in 2016. Since 2017, observed spawn mileage within Tenakee Inlet had shown a continual decline: 1.4 nmi in 2018, 0.5 nmi in 2019, 0.15 nmi in 2020, and none observed in 2021 or 2022. In 2023, 1.4 nmi of spawn was observed inside the entrance of Tenakee Inlet in the vicinity of South Passage Point and in 2024, 4.7 nmi of spawn was observed between Crab Bay and South Passage Point.

The spawning dates in 2024 were April 16–21, which is substantially earlier than the long-term average date of first spawn of April 27. Spawning in Tenakee Inlet has generally occurred between the last week in April and the first week in May (Tables 5 and 7). Historically, herring have spawned primarily in Tenakee Inlet along the southern shoreline between Saltery Bay and South Passage Point, with the core areas centered east and west of the Kadashan River flats. Herring spawn has also been documented intermittently along the Chatham Strait shoreline from South Passage Point south to Point Hayes, and more recently extending south to Lull Point on Catherine Island and west of Point Hayes along the northern entrance of Peril Strait. Most spawn observed in the last 5 seasons (2020–2024) has been along the Chatham Strait shoreline. The 2024 Chatham Strait–Peril Strait spawning event was similar to 2023 with nearly 6.0 nmi of spawn observed outside of Tenakee Inlet with most substantial spawning events occurring in the vicinity of Point Hayes.

No commercial herring fisheries will occur in Tenakee Inlet during the 2024–2025 season. Aerial surveys will begin in mid-April of 2025.

HOONAH SOUND (SECTION 13-C)

ADF&G began monitoring the Hoonah Sound herring stock in 1971. From 1971 through 2015, an average of 7.6 nmi of herring spawn was observed with an estimated average spawning biomass of 4,118 tons (Table 9). From the emergence of the Hoonah Sound SOK fishery in 1990, through 2015, the stock had averaged 10.3 nmi of spawn and 6,149 tons of estimated spawning biomass (Table 9). The highest estimated herring spawning biomass of 19,975 tons was observed in 2008. No spawn has been observed in Hoonah Sound since 2015.

When the SOK fishery was established in Hoonah Sound in 1990, the minimum threshold at which a fishery could occur was reduced from 2,000 tons to 1,000 tons. In 2015, to be more consistent with similar sized stocks around the region, the threshold in Hoonah Sound was increased to 2,000 tons. A summary of historical herring spawn timing and herring SOK harvest in Hoonah Sound is available in Tables 8 and 10. There has been no Hoonah Sound SOK fishery since a limited openpound fishery occurred in 2013.

No estimate of spawning biomass has been generated since 2015. The biomass in this area sharply decreased from the third largest stock in the region in 2008 to one of the smallest in 2012 and, based on aerial surveys, appears to have remained at a very low level since that time. There was no forecast for the 2024–2025 season; therefore, no commercial fishery will occur in Hoonah Sound in 2025.

CALENDAR OF EVENTS

The following is a calendar of events to be considered by pound operators for the 2025 herring season.

By February 28	2025 Southeast Alaska Herring SOK Pound Fishery Management Plan will be available at all Southeast Alaska area offices.
March 4	Advisory Announcement of the 2025 Ernest Sound, Hoonah Sound, and Tenakee Inlet closures, and the final 2025 Craig/Klawock GHL.
March 17	The Craig/Klawock fishery will open to seining of herring for placement in pounds effective 12:00 noon.

May 31 Pounds and all associated equipment in support of the fishery must be completely removed from the waters of the herring pound fishing area in Section 3-B. This includes the area covered by extreme high tide.

REGULATIONS

GENERAL SPAWN-ON-KELP REGULATIONS

The regulatory framework for the SOK fishery is found in 5 AAC 27.185. *Management plan for herring spawn on kelp in pounds fisheries in Sections 3-B, 12-A, and 13-C, and District 7.*

Placement and Release of Herring in Pounds

Herring may be placed in or added to a pound for 4 days, starting with the initial placement of herring in a pound. After 11:59 p.m. on the fourth day, no additional herring may be added to the pound (5 AAC 27.185 [q]). All herring in the pound must be released by 12:00 noon on the seventh day after the initial placement of herring in a pound (5 AAC 27.185 [s]). Under 5 AAC 27.185 (s), the first day is defined as the day herring are first placed into a pound. Once herring have been released or SOK product has been harvested, no additional herring or kelp may be introduced into a pound (5 AAC 27.185 [q]). Permit holders must take responsibility to ensure that when adding herring to a pound, herring are not at the same time swimming out of the pound, as this would constitute a violation of 5 AAC 27.185 (q). When releasing herring, at least one full side of the pound's webbing must be lowered a minimum of 2 feet below the surface of the water (5 AAC 27.185 [s]).

Post-Harvest Requirements

After a permit holder releases herring and harvests product from the pound, the permit holder must maintain the webbing in place for at least 4 weeks. To optimize hatching success, the permit holder must position egg-covered webbing in the original size and configuration of the pound structure with adequate water circulation on all sides. The webbing support system must be above the surface of the water and clearly marked as per 5 AAC 27.185 (k).

Harvest and Production

Each permit holder's SOK blades must remain separate from other permit holder's SOK blades until after processing and grading is completed. Permit holders will be allowed to harvest all SOK produced in their pounds. A permit holder's fish ticket must report only the SOK they harvested from their pound with the amount of blades used on the fish ticket. Each permit holder fishing a jointly operated pound shall be issued a fish ticket and the sum of the weights of those tickets shall equal the total weight of product produced in the jointly operated pound. For any product that has been delivered on the grounds to a licensed processor, the processor (not the permit holder) will be required to contact ADF&G with delivery weight for each landing on board.

REQUIREMENTS FOR BUYERS

Reporting requirements for buyers and processors of SOK product from Southeast Alaska SOK fisheries can be found in 5 AAC 27.187 *Buyer and processors reporting requirements for spawn on kelp in pounds for the Southeastern Alaska Area*. Buyers, processors, and permit holders should read and become familiar with these reporting requirements.

CLOSURE OF SOME POUND TYPES

ADF&G may close fishing for some pound types, if necessary, to avoid exceeding the GHL. In years when the GHL is low, instead of closing the fishery, ADF&G may instead close the fishery to certain pound types, such as single or double closed. There will be no restrictions on pound types for the 2025 Craig/Klawock SOK fishery.

OTHER REGULATIONS

Additional regulations pertaining to the pound fisheries can be found in the 2024–2025 Statewide Commercial Herring Fishing Regulations booklet under Chapter 27, Article 4, Southeast Alaska area under the following sections: 5 AAC 27.110 *Fishing seasons for Southeastern Alaska Area* (f), 5 AAC 27.130 *Lawful gear for Southeastern Alaska Area* (d), and (e); 5 AAC 27.185 *Management plan for herring spawn on kelp in pounds fisheries* (a) through (dd); 5 AAC 27.187 *Buyer and processors reporting requirements for spawn on kelp in pounds for the Southeastern Alaska Area*; harvesting requirements for *Macrocystis* kelp in 5 AAC 37.100 *Permits*; and 5 AAC 37.300 *Harvesting requirements for macrocystis kelp*.

Under 5 AAC 27.185 (w) all pounds and associated equipment used in these fisheries must be removed from the water by a specific date for a specific area. In Section 3-B, a permit holder shall completely remove all pounds and associated equipment from the waters by May 31. ADF&G and the Alaska Wildlife Troopers are advising permit holders that any pounds, nets, buoys, lines, and anchors left on the grounds will be removed and impounded or destroyed.

It is the responsibility of permit holders to carefully review and follow these regulations.

HARVEST AND ALLOCATION OF KELP FOR 2025

A permit issued by ADF&G is required to harvest kelp for use in pounds (5 AAC 37.900). Kelp harvest permits may be obtained from local ADF&G offices. Kelp blades will be allocated equally among permit holders fishing the same type of gear. The amount of kelp allowed for harvest for each permit holder is based on the kelp allocation table as indicated under regulation 5 AAC 27.185 (c) plus an allowance for breakage and loss during transport.

SECTION 3-B (CRAIG/KLAWOCK)

- Single permit closed pound—600 blades of *Macrocystis* kelp
- Double permit closed pound—900 blades of *Macrocystis* kelp
- Triple permit closed pound—1,000 blades of *Macrocystis* kelp

- Quadruple permit or more closed pound—1,000 blades of *Macrocystis* kelp
- Single permit open pound—3,000 blades of Macrocystis kelp
- Multiple permit open pounds—9,000 blades of *Macrocystis* kelp

FISHERY CONDUCT AND MANAGEMENT

The Craig/Klawock herring pound fishery will be the only SOK fishery in Southeast Alaska in 2025.

ADF&G will be closely monitoring herring activity using vessel and aerial surveys as well as communication with the harvesters. Results of aerial surveys will be announced by department advisory announcements on the day that an aerial survey occurs and will usually be updated on the grounds on Marine VHF Channel 10. Herring survey data collected by ADF&G from aerial and boat surveys will be available online through the interactive herring spawn maps located on the ADF&G's website¹. Regulation 5 AAC 27.185 (q) prevents a permit holder from releasing any herring from their pound when they are adding fresh herring.

To avoid mortality, the transport of herring to the pound site should be done with the pound itself or in a net pen that can be pushed or towed. Net pens used only for transporting herring must be marked *Tow Pound* and include the CFEC permit number of a participating permit holder. Transporting with a purse seine is discouraged except for very short distances. Pound operators should slowly push pounds or tow the pound alongside to avoid propellor wash and prevent crushing herring against the net. Permit holders are asked to avoid making and holding large sets to avoid herring mortality and stress.

Although regulations determine the maximum allowable number of kelp blades that can be harvested and placed in each permit holder's pound, permit holders are encouraged to fish the number of blades which will provide the maximum overall quality and value of their product rather than simply to fish the total amount allowed. Other measures have successfully been used in the fishery that may be considered when trying to maximize SOK quality and value including the following:

- 1) Pound nets may be shaped with internal frames to provide the full net volume.
- 2) Kelp depths in the pound may be matched with the depth of active spawning by testing spawn deposition with a weighted string.
- 3) Fishing and transferring herring to pounds should only occur once herring are fully mature.
- 4) Top-off sets may be added over a 4-day period once herring have been initially introduced.
- 5) Herring density in the pound should be closely monitored since successful spawning is inhibited by excessive crowding.
- 6) Web depth may be adjusted (consistent with specifications under 5 AAC 27.130 (e)(1)(C) or 5 AAC 27.185 (cc) to provide good water exchange.

¹ Interactive Maps. Commercial Herring Fisheries: Southeast Alaska and Yakutat. 2021–. Alaska Department of Fish and Game. Juneau, AK. http://www.adfg.alaska.gov/index.cfm?adfg=commercialbyareasoutheast.herring#maps.

OPEN WATERS

In Section 3-B, certain areas are open to the operation of pounds for taking of herring SOK and seines for taking of herring for placement in pounds. Those areas are shown in Figure 1, and include:

- Gulf of Esquibel
- San Alberto Bay
- those waters of Shinaku Inlet and San Christoval Channel south of the latitude of the northernmost tip of Saint Philip Island at 55°39.31' N lat, 133°25.12' W long, east of a line from the northernmost tip of Saint Philip Island to the northernmost tip of Point Garcia at 55°33.65' N lat, 133°26.47' W long, and north of a line from Entrance Point at 55°31.16' N lat, 133°08.69' W long to the southernmost tip of Clam Island at 55°30.98' N lat, 133°09.45' W long to the southernmost tip of Fern Point 55°30.05' N lat, 133°16.97' W long.

CLOSED WATERS

In Section 3-B, certain areas are closed to the operation of pounds for taking of herring SOK and seines for taking of herring for placement in pounds. Those areas are shown in Figure 1, and include:

- Klawock Inlet and Big Salt Lake east of the longitude of the northern point of Wadleigh Island at 133°07.57' W long.
- those waters of San Christoval Channel in the main channel enclosed by a line from 55°35.62' N lat, 133°20.00' W long to 55°35.17' N lat, 133°20.00' W long to 55°33.37' N lat, 133°17.52' W long to 55°33.50' N lat, 133°17.28' W long.
- those waters of Fish Egg and Ballenas Islands south of 55°31.00' N lat, north of the southernmost tip of Cape Suspiro at 55°27.47' N lat, 133°08.40' W long, and east of the longitude of Ballena Island Shoal Light at 133°13.25' W long.

OTHER AGENCY REQUIREMENTS

Prospective pound operators are advised to consider other agency requirements for constructing and operating pounds in Craig/Klawock, Ernest Sound, Tenakee Inlet, and Hoonah Sound. Pound operators are urged to contact the Alaska Department of Natural Resources (ADNR), the U.S. Forest Service (USFS), the U.S. National Marine Fisheries Service (NMFS), and the U.S. Coast Guard (USCG) to determine other regulations and requirements. Phone numbers for those agencies are listed below.

ALASKA DEPARTMENT OF NATURAL RESOURCES

ADNR manages the use of tidelands and submerged lands seaward of mean high water. Contact ADNR at (907) 465-3400.

U.S. FOREST SERVICE

The U.S. Forest Service has jurisdiction over and manages most of the lands above mean high tide. People who plan to use National Forest land in connection with the fishery must apply for a special use permit from USFS prior to any occupancy. Special use permit applications are available from local USFS offices. Completed applications should be submitted well in advance of operations to ensure that a permit is received in time for the fishery. Examples of use needing a permit include (but are not limited to) camping or storage of gear on National Forest land in conjunction with the commercial fishery. Contact USFS at (907) 225-3101.

U.S. NATIONAL MARINE FISHERIES SERVICE

Marine mammal interactions should be reported to NMFS at (907) 586-7221.

U.S. COAST GUARD

Structures such as floating fish pens are subject to the requirements of the Code of Federal Regulations, Title 33, Part 64. This regulation requires an owner to apply for a USCG permit and to install and maintain a light or other private aid to navigation if the USCG determines it to be necessary to protect maritime navigation.

Herring pounds used in the SOK pound fishery do not require permits for private aids to navigation at this time, provided the owners:

- Place 2 signs on opposite corners of the structure. These signs will be worded "Danger, Fish Pens" (Figure 3).
- Place a single, all-points white light on one corner of structures less than 400 square feet in size.
- Place a single, all-points white light on every corner of structures larger than 400 square feet in size.
- Anchor fish pens within the boundary areas specified in ADF&G regulation 5 AAC 27.185 (f) (Figures 1 and 2).

If all these conditions are not met, the permit holder must apply to the USCG for an individual *Private Aids to Navigation Permit*. If you have questions, call the USCG Aids to Navigation office, at (907) 463-2254.

LIST OF CONTACTS

The following ADF&G, Division of Commercial Fisheries personnel may be contacted regarding this management plan:

Bo Meredith, Justin Breese, and Whitney Crittenden	2030 Sea Level Dr., Suite 205			
Area Management Biologists	Ketchikan, Alaska 99901			
Ketchikan Area Office	(907) 255-5195			
Aaron Dupuis and Anthony Walloch	304 Lake St. Rm. 103			
Area Management Biologists	Sitka, Alaska 99835			
Sitka Area Office	(907) 747-6688			
Scott Forbes and Raymond Vinzant	P.O. Box 110024			
Area Management Biologists	Juneau, Alaska 99811			
Douglas Regional Office	(907) 465-4250			
Katie Taylor, Tom Kowalske, and Emily Klosterman	P.O. Box 667			
Area Management Biologists	Petersburg, Alaska 99833			
Petersburg Area Office	(907) 772-3801			
Kyle Hebert	P.O. Box 110024			
Herring Research Biologist	Douglas, Alaska 99811			
Douglas Regional Office	(907) 465-4250			
Troy Thynes	P.O. Box 667			
Region I Management Coordinator	Petersburg, Alaska 99833			
Petersburg Area Office	(907) 772-3801			
Anne Reynolds Manney	2030 Sea Level Dr., Suite 205			
Region I Supervisor	Ketchikan, Alaska 99901			
Ketchikan Area Office	(907) 255-5195			

TABLES AND FIGURES

			Total GHL ^c				
		Forecasted pre-	bait and		Bait	Number	
	Miles of	fishery	SOK^d	Bait GHL	harvest	of	Exvessel
Season ^a	spawn	biomass ^b	(tons)	(tons)	(tons)	permits	value
1990/91	22.0	18,350	—	2,841	3,272	27	\$981,600
1991/92	23.0	17,800	2,684	2,281	2,295	28	\$619,650
1992/93°	8.4	12,350	1,602	1,362	623	10	\$150,960
1993/94	8.0	7,996	895	760	636	6	\$187,578
1994/95	5.5	6,778	725	617	***	***	***
1995/96	9.9	6,262	658	558	***	***	***
1996/97	13.2	6,755	715	615	517	4	\$137,788
$1997/98^{\rm f}$	11.0	7,018	755	455	***	***	***
1998/99	15.4	6,951	750	450	***	***	***
1999/00	12.9	6,013	626	376	***	***	***
2000/01	16.7	9,091	1,058	635	***	***	***
2001/02	18.0	8,387	952	571	***	***	***
2002/03	11.2	6,045	630	378	***	***	***
2003/04	12.0	13,204	1,754	1,052	***	***	***
2004/05	18.0	15,577	2,217	1,330	553	3	\$199,012
2005/06	8.2	14,262	1,955	1,173	689	3	\$247,934
2006/07	22.3	13,768	1,860	1,116	576	3	\$139,000
2007/08	11.0	14,213	1,945	1,167	565	3	\$133,300
2008/09	17.0	14,213	1,945	1,167	142	3	\$51,304
2009/10	18.7	14,870	2,074	1,244	***	***	***
2010/11	14.8	17,886	2,710	1,140	***	***	***
2011/12	14.9	34,235	6,847	4,060	309	3	\$113,784
2012/13	15.3	23,391	4,060	2,436	321	3	\$116,000
2013/14	13.6	26,085	4,808	2,884	***	***	***
2014/15	11.5	15,803	2,362	1,367	964	3	\$396,205
2015/16	12.3	12,303	1,590	954	898	3	\$369,660
2016/17	22.8	7,833	872	523	527	3	\$210,800
2017/18	17.3	16,083	2,312	1,387	527	5	\$310,000
2018/19	28.9	22,810	3,906	2,344	710	6	\$301,750
2019/20	56.1	55,072	11,014	6,608	995	4	\$410,000
2020/21	34.2	97,282	19,456	11,674	540	5	\$225,000
2021/22	36.4	63,250	12,650	7,590	398	2	\$168,000
2022/23	29.4	38,804	7,761	4,656	742	3	\$296,640
2023/24	23.8	33,580	6,716	4,030	***	***	***
2024/25	ND	30,182	6,036	3,622	ND	ND	ND
Average	17.4	20,147	3,180	1,963	493	4	\$251,405

Table 1.-Craig/Klawock herring stock size and fishery summary, 1990-2024.

Note: *** indicates data is confidential. ND indicates data were still forthcoming at time of analysis. En dashes indicate no data. ^a Spawn year is second year of regulatory season listed in the adjacent year column.

^b Forecasted pre-fishery biomass values were estimated with hydroacoustic surveys for 86/87, spawn deposition surveys

for 87/88 to 92/93, and age-structured models for 93/94 to 18/19.

^c Guideline harvest limit (GHL).

^d Spawn on kelp (SOK).

^e First year bait quota was split with pound fishery 85%:15%.

^f Herring allocation changed to 60% for the winter food and bait fishery and 40% to the pound fishery.

υ	01	Ĩ	, ,	,		
	2010	2011	2012	2013	2014	2015
Herring SOK quota (tons) ^a	1,953	2,402	6,538	3,739	4,544	1,308
Total harvest SOK (tons)	116.7	70	98.1	137.7	***	***
				\$3,099,00		
Exvessel value	\$884,715	\$728,147	\$2,099,002	2	***	***
Average price/lb	\$3.80	\$5.13	\$10.69	\$12.00	***	***
Average income	\$8,268	\$14,003	\$32,795	\$23,656	***	***
Number of pounds	63	34	35	80	75	76
Number of landings	107	52	64	131	136	135
Blade allocation	b	b	b	b	b	b
Total kelp harvest (tons)	8.2	4.6	5.3	9.3	19.2	19.2
Herring spawning dates	4/5-4/14	4/1-4/7	4/3-4/8	3/31-4/3	4/1-4/5	3/27-4/1
Miles of spawn	18.7	14.8	14.9	15.3	13.6	11.5
Forecasted pre-fishery						
biomass (tons)	14,870	17,886	34,235	23,391	26,085	15,803
X /	2016	2017	2018	2019	2020	2021
Herring SOK quota (tons) ^a	692	349	1,602	2,895	10,119	18,916
Total harvest SOK (tons)	***	69.9	205.3	202.4	283.9	262.6
		0,1,1	20010	\$3,300,00		_0_0
Exvessel value	***	\$932,917	\$3,262,900	0	\$3,256,000	\$2,150,000
Average price/lb	***	\$6.68	\$7.95	\$8.15	\$5.73	\$4.77
Average income	***	\$8,042	\$24,533	\$23,358	\$21,853	\$15,468
Number of pounds	46	19	66	73	75	80
Number of landings	133	116	133	141	149	158
Blade allocation	c	d	e	e	e	e
Total kelp harvest (tons)	9.2	10.2 3/24-4/2 &	9.4	9.8	11.4	12.5
Herring spawning dates	3/25-3/31	4/6-4/8	3/29-4/4	3/27-4/6	4/3-4/13	4/5-4/14
Miles of spawn	12.3	22.8	17.3	28.9	56.1	34.2
Forecasted pre-fishery	12.5	22.0	17.5	20.9	50.1	51.2
biomass (tons)	12,303	7,833	16,039	22,810	86,087	97,282
	2022	2023	2024	22,010	00,007	97,202
Herring SOK quota (tons) ^a	12,252	7,078	6,450	_		
Total harvest SOK (tons)	12,252	127.9	0,430 92.8			
Exvessel value	\$2,437,880	\$1,950,471	92.8 \$2,117,340			
Average price/lb	\$2,437,880 \$6.30	\$1,930,471 \$7.65	\$2,117,540 \$11.23			
Average income	\$0.30 \$20,063	\$7.03 \$25,006	\$11.25			
Number of pounds	\$20,005 60	\$23,000 41	\$33,083 33			
	131	87	55 64			
Number of landings Blade allocation	131 e	8 / e	04 e			
Total kelp harvest (tons)		8.6				
i otar keip narvest (tons)	9.1	0.0	2/21 2/27			
II	A/5 A/1 C	A/1 A/11	3/21-3/27,			
Herring spawning dates	4/5-4/16	4/1-4/11	4/4 22.8			
Miles of spawn	36.4	29.4	23.8			
Forecasted pre-fishery	(2.25)	20.004	22.590			
biomass (tons)	63,250	38,804	33,580	_		

Table 2.-Craig/Klawock herring spawn-on-kelp (SOK) fishery summary, 2010-2024.

Note: *** indicates data is confidential.

^a The quota was updated to include the total SOK quota which includes the carry-over from unharvested winter bait.

^b 600 blades for a single closed pound, 750 blades for a double closed pound, 1,125 blades for a triple closed pound.

^c 300 blades for a single closed pound, 400 blades for a double closed pound, 700 blades for a triple closed pound.

^d 500 blades per permit with 6 permit holders per pound structure.

^e 600 blades for a single closed pound, 900 blades for a double closed pound, 1,000 blades for a triple closed pound.




Table 3.-Craig and Ernest Sound herring spawning dates comparison, 2010-2024

Note: Black bars indicate spawning dates for Craig and gray bars indicate spawning dates for Ernest Sound.

Season	Date of first spawn ^a	Nautical miles of spawn ^b	Forecast for GHL determination ^{c,d} (tons)	Spawning biomass (tons) ^e	GHL (tons) ^f	Bait harvest (tons) ^g	SOK harvest (lbs)	Sac Roe harvest (tons)	Remaining GHL (tons)
1969/70	_	_	—	_	_	17	_	_	_
1970/71	_	3	_	13,100	_	206	_	_	_
1971/72	_	_	13,100	3,650	_	967	_	_	_
1972/73	_	_	3,650	450	_	775	_	_	_
1973/74	_	_	450	400	_	535	_	_	_
1974/75	_	_	400	2,900	_	593	_	_	_
1975/76	_	3	2,900	4,350	580	708	_	_	_
1976/77 ^g	_	3	4,350	3,035	870	901	_	49	0
1977/78	5/3	_	3,035	1,505	455	340	_	_	0
1978/79	4/16	2.6	1,505	255	_	_	_	_	115
1979/80	5/2	4	255	500	_	_	_	_	_
1980/81	_	3.5	500	410	_	_	_	_	_
1981/82	_	_	410	160	_	_	_	_	_
1982/83	_	_	160	1,640	_	_	_	_	_
1983/84	4/11	_	1,640	1,000	_	_	_	_	_
1984/85	_	4.5	1,000	1,000	_	_	_	_	_
1985/86	_	_	1,000	1,000	_	_	_	_	_
1986/87	_	1	1,000	_	_	_	_	_	_
1987/88	4/21	2	_	_	_	_	_	_	_
1988/89	4/17	2.4	_	500	_	_	_	_	_
1989/90	_	2.1	500	1,000	_	_	_	_	_
1990/91	_	_	1,000	3,000	_	_	_	_	_
1991/92	4/16	9.1	3,000	2,650	_	_	_	_	_
1992/93	4/23	9	2,650	684	200	8	_	_	192
1993/94	4/26	8.4	684	2,544	0	_	_	_	_
1994/95	4/25	6.5	2,544	2,470	255	111	_	_	144
1995/96	4/16	6.9	2,744	2,665	280	220	_	_	60
1996/97	4/16	0	4,852	0	377	6	_	_	371
1997/98	4/9	11.8	_	5,998	0	_	_	_	_
1998/99	4/5	1.8	5,381	No survey	662	96	_	_	566
1999/00	4/8	9.1	_	920	0	_	_	_	_
2000/01	4/11	6.9	_	2,052	0	_	_	_	_
2001/02	4/15	4.8	1,653	2,406	0	_	_	_	_
2002/03	4/10	8.5	2,407	5,509	0	_	_	_	_
2003/04	4/11	7.1	6,592	2,413	875	44	112,286	_	831
2004/05	4/22	10.1	1,906	3,268	0	_	_	_	_
2005/06	4/6	7.9	2,284	2,538	0	_	_	_	_

Table 4.–Ernest Sound herring stock and fishery summary, 1969–2024.

-continued-

Season	Date of first spawn ^a	Nautical miles of spawn ^b	Forecast for GHL determination ^o (tons)	Spawning ^{,d} biomass (tons)°	GHL (tons) ^f	Bait harvest (tons) ^g	SOK harvest (lbs)	Sac Roe harvest (tons)	Remaining GHL (tons)
2006/07	4/19	11.3	1,955	7,353	0				
2007/08	4/20	15.4	9,060	4,846	1382	***	19,650	_	>700
2008/09	4/17	6.6	4,545	2,862	529	***	4,911	_	100-299
2009/10	4/14	7.8	2,879	3,523	297	***	_	_	<50
2010/11	4/17	8.1	5,080	2,559	476	***	0	_	100-299
2011/12	4/16	8.9	2,682	3,193	272	***	_	_	<50
2012/13	4/16	9.7	3,509	7,556	379	***	129,265	_	100-299
2013/14	4/14	3.7	7,613	2,631	1073	***	***	_	>700
2014/15	4/8	5.5	1,991	562	0	_	_	_	_
2015/16	4/10	4.4	1,207	346	0	_	_	_	_
2016/17	4/14	4.4	_	_	_	_	_	_	_
2017/18	4/24	3.5	_	_	_	_	_	_	_
2018/19	4/14	1	_	_	_	_	_	_	_
2019/20	4/19	0.3	_	_	_	_	_	_	_
2020/21	4/15	0.5	_	_	_	_	_	_	_
2021/22	4/14	2.6	_	_	_	_	_	_	_
2022/23	4/23	2.0	_	_	_	_	_	_	_
2023/24	4/9	3.8	_	_	_	_	_	_	_
Recent 10-yr Average	4/15	2.8	1,599	454	0	_	_	_	_
Long- term Average	4/16	5.5	2,852	2,591	332	44	***	_	253

Note: En dashes indicate no data. *** indicates data is confidential. Spawn on kelp is abbreviated to SOK and guideline harvest limit is abbreviated to GHL.

^a Since the 1997/1998 season, the first spawn and the major spawn have been within 5 days of each other.

^b 1996/1997– No spawn deposition survey conducted; 7.5 miles of spawn observed along Ship Island;

1998/1999- No spawn deposition survey conducted; 1.8 miles of spawn observed

^c From 1971/1972 through 1984/1985 season forecasts were based on computer generated estimates derived using hydroacoustic techniques; from 1985/1986 through 1991/1992 forecasts were based on visual estimates; from 1992/1993 through 1994/1995 forecasts were based on spawn deposition estimates; from 1995/1996 through 2006/2007 biomass accounting was used.

^d Beginning with the 1976/1977 season fishery threshold established at 2,500 tons.

^e From1969/1970 through 1983/1984 biomass estimates were -computer generated estimates derived using hydroacoustic techniques; from1984/1985 through 1990/1991 estimates were based on visual observations and from1991/1992 through 2005/2006 were spawn deposition estimates. 1975/1976 and 1976/1977 GHLs are based upon 20% of the hydroacoustic estimate. 1977/1978 GHL is based upon 15% of the hydroacoustic estimate.

^f 2003/2004 GHL includes 90 tons rolled over from the bait pound fishery.

^g 1973/1974,1974/1975, 1976/1977 seasons also include harvests from Fools and Menefee Inlets. Does not include harvests from statistical area 107-40.

Statistic	2004	2008	2009
GHL (tons)	875	1,382	529
GHL available for SOK (tons)	775	***	***
SOK harvest (tons)	56.1	9.8	2.5
Exvessel value	\$514,900	\$131,100	\$18,400
Average price/lb	\$4.59	\$6.87	\$3.67
Average income/permit	\$8,046	\$10,091	\$4,590
Number of permits	64	13	4
Number of pounds	51/6/0/1ª	1/6/0/0 ^a	0/2/0/0 ^a
Number of landings	64	13	4
Kelp allocation (blades)	b	c	d
Kelp blade harvest (lbs)	4,600	29,400	2,400
Fishery open-closed	4/1-5/10	4/1-5/10	4/1-5/10
Fishing occurred	4/10-4/17	4/16/-4/24	4/17-4/21
Harvest occurred	4/14-1/17	24-Apr	21-Apr
	2011	2013	2014
GHL (tons)	476	379	1,073
GHL available for SOK (tons)	***	***	***
SOK harvest (tons)	0	64.3	***
Exvessel value	\$0	\$1,574,700	***
Average price/lb	\$0	\$12.25	***
Average income/permit	\$0	\$19,441	***
Number of permits	0	81	129
Number of pounds	0	1/5/1/22/0 ^e	25/52/0ª
Number of landings	0	81	129
Kelp allocation (blades)	d	d	c
Kelp blade harvest (lbs)	0	6,400	29,000
Fishery open-closed	4/1-5/10	4/1-5/10	4/1-5/10
Fishing occurred	NA	4/10-4/16	4/12-4/20
Harvest occurred	NA	4/15-4/16	4/18-4/20

Table 5.-Ernest Sound spawn-on-kelp (SOK) fishery summary, 2004, 2008, 2009, 2011, 2013, and 2014.

Note: *** indicates data is confidential. NA indicates no harvest. Guideline harvest limit is abbreviated to GHL.

^a Single/double/triple/open.

^b 1,000 blades single closed/1,000 blades double closed/1,000 blades triple closed/2,500 blades single-open/7,500 multiple permit open.

^c 1,000 blades single closed/2,000 blades double closed/1,000 blades triple closed/2,500 blades single-open/7,500 multiple permit open.

^d 200 blades single closed/400 blades double closed/500 blades triple closed/1,500 blades single-open/4,500 multiple permit open.

^e Single/double/combined double/triple/open.

Season	Major spawning dates	Nautical miles of spawn	Spawning biomass ^a (tons)	Food/bait GHL (tons)	Food/bait harvest (tons)
1978/79	5/9-5/11	3.3	2,500	200	0
1979/80	4/28-5/2	3.9	4,485	400	504
1980/81	4/27-5/5	9.3	7,500	750	847
1981/82	4/25-5/7	11.1	6,650	650	687
1982/83	4/25-5/6	13.1	8,870	875	749
1983/84	4/20-4/26	8.3	12,100	850	619
1984/85	4/24-5/1	9.9	11,000	1,400	1,406
1985/86	4/27-5/1	8.3	12,500	1,700	2,040
1986/87	4/22-4/30	7.9	6,600	800	1,275
1987/88	4/22-4/27	9.1	6,000	1,450	1,577
1988/89	4/26-4/29	10.3	5,360	720	655
1989/90	4/25-5/6	2.9	2000	650	595
1990/91	4/25-5/4	2.1	400	_	_
1991/92	5/5	trace	200	_	_
1992/93	4/21-4/23	6.4	904	_	_
1993/94	4/24-4/26	0.25	400	_	_
1994/95	4/26	0.05	200	_	_
1995/96	5/4-5/14	18.1	4,560	_	_
1996/97	4/26-5/7	14.4	9,926	300	98
1997/98	4/24-4/29	12.4	10,419	825	586
1998/99	4/25-4/28	11	11,049	1,023	835
1999/00	4/26-5/3	13.8	9,425	542	494
2000/01	4/21-5/1	12.2	7,576	906	775
2001/02	4/23-4/27	15.4	4,084	840	355
2002/03	4/25-4/28	12.2	3,529	528	328
2003/04	4/28-5/3	13	4,728	399	***
2004/05	4/26-5/2	8.9	3,036	476	0
2005/06	5/2-5/6	5.9	5,110	_	_
2006/07	4/23-4/26	4.4	3,346	_	_
2007/08	4/30, 5/7–5/8	11.4	11,252	_	_
2008/09	4/25-4/26, 4/29-4/30	6.9	5,283	875	254
2009/10	5/7-5/9	2.7	1,437	583	***
2010/11	5/9	1	_	_	_
2011/12	4/20-4/23	4.6	5,119	_	_
2012/13	5/7-5/10	5.4	4,936	_	_

Table 6.–Tenakee Inlet herring stock and fishery summary, 1978–2024.

-continued-

Season	Major spawning dates	Nautical miles of spawn	Spawning biomass ^a (tons)	Food/bait GHL (tons)	Food/bait harvest (tons)
2013/14	4/29	2	927	557	***
2014/15	4/22-4/25	2.3	2,223	_	_
2015/16	-	0	_	_	_
2016/17	5/13-5/16	2.1	_	_	_
2017/18	5/7-5/9	1.4	_	_	_
2018/19	5/12-5/14	0.5	_	_	_
2019/20	5/5-5/6	1.9	_	_	_
2020/21	4/20-4/21	1.7	_	_	_
2021/22	4/28	spot	_	_	_
2022/23	4/24, 5/3–5/4	7.6	_	_	_
2023/24	4/17-4/20	10.6	_	_	_

Table 6.–Page 2 of 2.

Note: En dashes indicate no data and *** indicates data is confidential. Guideline harvest limit is abbreviated to GHL.

^a Spawning biomass estimates were calculated from hydro-acoustical surveys from 1979 through 1986, and from spawn deposition surveys from 1987 through 2015—bolded values were derived from ASA models.

	\mathcal{O} 1		
Statistic	2003	2004	2005
GHL (tons)	528	399	476
GHL available for SOK	180	***	476
SOK harvest (tons)	47.6	100.7	101.4
Exvessel value	\$580,500	\$981,500	\$512,900
Average price/lb	\$6.10	\$4.68	\$2.53
Average income/permit	\$10,560	\$11,680	\$5,640
Number of permits	55	85	98
Number of pounds	1/15/8/0ª	1/32/6/2/2 ^b	1/29/13/3ª
Number permits landing	55	85	91
Kelp allocation (blades)	200/400/550/0ª	300/500/500/2000 ^a	300/500/500/2000 ^a
Kelp blade harvest (lbs)	35,375	39,000	53,850
Fishery open-closed	4/6-5/6	4/6-5/6	4/6-5/5
Fishing occurred	4/25-4/28	4/28-5/1	4/27-4/30
Harvest occurred	4/30-5/4	5/3-5/6	5/2-5/4
Statistic	2009	2014	
GHL (tons)	875	557	
GHL available for SOK	621	***	
SOK harvest (tons)	64.1	84.4	
Exvessel value	\$558,900	\$1,155,300	
Average price/lb	\$4.36	\$6.85	
Average income/permit	\$6,500	\$16,270	
Number of permits	86	78	
Number of pounds	$11/27/7/0^{a}$	2/25/5/1°	
Number permits landing	86	71	
Kelp allocation (blades)	400/500/500/0ª	300/500/500/0 ^a	
Kelp blade harvest (lbs)	42,600	41,250	
Fishery open-closed	4/6-5/5	4/6-5/4	
Fishing occurred	4/28-5/1	4/27-5/1	
Harvest occurred	5/2-5/5	5/2-5/4	

Table 7.–Tenakee Inlet herring Spawn-on-kelp (SOK) fishery summary, 2003–2005, 2009, and 2014.

Note: *** indicates data is confidential. Guideline harvest limit is abbreviated to GHL.

^a Single/double/triple/test.

^b Single/double/triple/long line/test.

^c Single/double/triple/quadruple.

									April	1													М	av						
Year	Location	16	17	18	19	20	21	22			25	26	27	28	29	30	1	2	3 4	5	6	7 8	9	10	11	12	13	14	15	16
2003	Hoonah Sound				- /												-						-							
	Tenakee																													
2004	Hoonah Sound																													
	Tenakee																													
2005	Hoonah Sound																													
	Tenakee																													
2006	Hoonah Sound																													
	Tenakee											1																		
2007	Hoonah Sound																													
	Tenakee																													
2008	Hoonah Sound																					_								
	Tenakee																													
2009	Hoonah Sound																													
2010	Tenakee																													
2010	Hoonah Sound Tenakee																													
2011	Hoonah Sound																													
2011	Tenakee																													
2012	Hoonah Sound																													
2012	Tenakee																													
2013	Hoonah Sound							-																						
2015	Tenakee																													
2014	Hoonah Sound																													
2011	Tenakee																						-							
2015	Hoonah Sound																													
	Tenakee																													
2016^{*}	Hoonah Sound																													
	Tenakee																													
2017	Hoonah Sound																													
	Tenakee																													
2018	Hoonah Sound																													
2010	Tenakee																													
2019	Hoonah Sound																													
2020	Tenakee																								_					
2020	Hoonah Sound Tenakee																													
2021	Hoonah Sound																	- 1												
2021	Tenakee				1																									
2022	Hoonah Sound				_	_	_																							
2022	Tenakee																													
2023	Hoonah Sound																													
2025	Tenakee									_	_								_		1									
2024	Hoonah Sound																				_									
2021	Tenakee																													

Table 8.–Hoonah Sound and Tenakee Inlet herring spawn dates comparison, 2003–2024.

Note: Black bars indicate spawning dates for Hoonah Sound and gray bars indicate spawning dates for Tenakee Inlet. * indicates year when no spawn was observed in either stock

Year	Spawn dates	Nautical miles of spawn	Spawning biomass (tons)	SOK harvest (tons)
1971	5/10-5/17	2.5	833	-
1972	5/11-5/12	1.5	666	-
1973	_	1.0	333	-
1974	5/14	3.0	999	_
1975	_			-
1976	5/5	1.0	333	-
1977	_	3.5	1,166	-
1978	_	5.3	1,765	-
1979	_	0.5	167	-
1980	_	-	_	-
1981	4/30-5/01	2.3	750	-
1982	4/29-5/01	1.5	500	-
1983	5/1	1.0	333	-
1984	4/26-5/01	3.0	540	-
1985	5/01-5/03	3.5	1,166	-
1986	4/28-5/01	3.8	1,249	-
1987	4/28-5/02	3.8	740	-
1988	4/30-5/01	5.0	1,665	-
1989	4/16-4/20	17.0	4,000	-
1990	4/13-4/28	10.0	2,350	12.0
1991	4/19-4/24	8.7	2,175	13.6
1992	4/22-4/24	10.8	5,714	23.1
1993	4/27-4/29	5.7	1,099	14.8
1994	4/21-4/23	9.0	2,450	33.3
1995	4/20-4/21	4.5	274	28.7
1996	5/02-5/9	10.1	4,023	-
1997	4/25-4/28	14.5	5,884	64.3
1998	4/23-4/27	14.5	6,472	85.6
1999	4/27-5/1	13.8	4,426	93.8
2000	4/27-4/30	13.0	3,635	36.0
2001	4/27-5/1	13.7	8,538	66.2
2002	4/25-4/27	11.9	4,936	136.6
2003	4/23-4/27	16.7	9,423	141.6
2004	4/22-4/29	11.1	7,502	236.1
2005	4/18-4/25	10.3	6,924	181.7
2006	4/23-4/26	9.0	6,028	162.1
2007	4/46-5/2	16.5	10,946	159.4
2008	4/23-4/30	14.5	19,975	228.1
2009	4/22-4/27	10.3	15,829	234.7
2010	4/22 - 4/28	12.4	15,264	290.4
2011	4/20-5/5	12.6	14,215	193.7
2012	4/20-4/23	4.2	923	186.5
2013	5/7-5/9	2.4	412	-
2014	5/5-5/8	3.2	444	-
2015	4/24-5/6	4.2	22	_
Average	1971-2015	7.6	4,118	-
Average	1990-2015	10.3	6,149	119.2

Table 9.-Hoonah Sound herring spawning stock and spawn-on-kelp (SOK) fishery summary, 1971–2015.

Note: En dashes indicate no data. Shaded estimated escapements are based on average spawn density from the years 1989 to 2002; due to funding, comprehensive aerial surveys have not been conducted since 2016. There has been no herring spawn observed, nor a biomass estimate generated since 2015 in Hoonah Sound.

	2003	2004	2005	2006	2007
Herring GHL (tons)	427	1,207	728	669	681
Harvest (tons)	141.6	237.4	190.6	162.1	144.5
Exvessel value	\$1,922,500	\$2,071,347	\$1,117,568	\$1,943,422	\$4,491,070
Average price/lb	\$6.79	\$4.36	\$2.93	\$6.00	\$14.09
Average income	\$17,800	\$19,541	\$11,889	\$24,600	\$49,352
Number of pounds	$49/1/3^{a}$	92/12/2 ^b	81/5/3°	17/45 ^a	67/12 ^a
Number selling product	108	106	94	79	91
Kelp allocation (blades)	500/300/750ª	1,000/1,000/	1,000/1,000/	2,500/1,000/	2,500/1,000/
		3,000 ^b	1,500 ^c	1,500 ^a	1,500 ^a
Kelp blade harvest	60,301	126,000	118,450	136,698	122,565
Fishery open-closed	4/6-4/25	4/6-4/28	4/6-4/28	4/6-4/27	4/6-5/4
Fishing occurred	4/19-4/24	4/20-4/25	4/19-4/28	4/18-4/23	4/23-4/29
Harvest occurred	4/24-4/27	4/26-4/28	4/25-4/28	4/23-4/27	4/30-5/4
	2008	2009	2010	2011	2012
Herring GHL (tons)	2,238	2,238	3,182	3,015	2,139
Harvest (tons)	223	234.7	290.4	193.7	186.5
Exvessel value	\$5,115,459	\$2,332,514	\$2,580,517	\$1,820,952	\$4,033,078
Average price/lb	\$11.47	\$4.97	\$4.44	\$4.70	\$10.81
Average income	\$51,155	\$23,094	\$25,550	\$20,460	\$55,248
Number of pounds	98/3ª	99/4 ^a	97/2ª	85/4 ^a	83/4 ^a
Number selling product	100	101	101	89	73
Kelp allocation (blades)	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/
	1,500 ^a	1,500ª	1,500 ^a	1,500ª	1,500ª
Kelp blade harvest	201,262	196,492	178,898	169,922	155,104
Fishery open-closed	4/6-5/02	4/6-4/30	4/6-4/30	4/6-4/30	4/6-5/15
Fishing occurred	4/22-4/27	4/22-4/25	4/21-4/25	4/20-4/24	4/19-4/21
Harvest occurred	4/27-5/1	4/26-4/29	4/22-4/28	4/26-4/29	4/26-4/27
	2013				
Herring GHL (tons)	130				
Harvest (tons)	0				
Exvessel value	\$0				
Average price/lb	\$0				
Average income	\$0				
Number of pounds	3 open				
Number selling product	0				
Kelp allocation (blades)	600 open				
Kelp blade harvest	7,940				
Fishery open-closed	4/6-5/15				
Fishing occurred	NA				
Harvest occurred	NA				

Table 10.-Hoonah Sound herring spawn-on-kelp (SOK) detailed fishery summary, 2003-2013.

Note: NA indicates no harvest. Guideline harvest limit is abbreviated to GHL. Fisheries did not occur in 2014–2022 since the biomass forecast was either below the 1,000-ton threshold or no survey was conducted due to minimum herring spawn observed.

^a Double closed pounds/single closed pounds/triple closed pounds.

^b Double closed pounds/single closed pounds/open pounds.

^c Single permit closed pound/double permit closed pound/triple permit closed pounds.




Figure 1.-Craig/Klawock (Section 3-B) herring SOK fishery open area.




Figure 2.–Diagram of a herring pound showing 2 alternative methods of marking herring pounds.

Note: Regulations require vertical signs with the permit holder's first and last name and five-digit CFEC permit number (5 AAC 27.185(k)). Letters and numbers must be at least 6 inches high and at least one-half inch wide, must contrast with the background and must be always above the waterline.




Figure 3.–U.S. Coast Guard requirements for marking pounds.